


Application Note

Exosome isolation from Cerebrospinal Fluid (CSF)



Exosome isolation from CSF using Exo-spin™ exosome purification kit

Data Courtesy of Tânia Soares Martins *et al.* 2018 as published in PLoS One journal University of Aveiro, Portugal (Issued March, 18th 2019)

- **Experiment**
Isolate exosomes using Exo-spin™
- **Exosomes origin**
CSF
- **Initial sample volume**
5 ml per Exo-spin™ column
- **Elution sample volume**
200 µl per sample in PBS

Summary

Exosomes may be able to pass through the blood-brain barrier although the mechanism is not well-understood. This property makes exosome research attractive for areas such as biomarker discovery for neurodegenerative diseases and also drug delivery. To advance these areas of research, efficient and reproducible methods for isolating exosomes from small volumes of CSF need to be developed.


In this study, the Exo-spin™ exosome isolation system is compared to two precipitation methods using 5 ml of CSF as the starting sample. The Exo-spin™ method has been identified as superior. Among other factors, yield, purity, as well as structural integrity of the generated samples have been analysed as part of this comparison. Nanoparticle Tracking Analysis (NTA), exosome protein to particle ratio, Western Blot and Transmission Electron Microscopy (TEM) were used to generate comparative data.

Key Features of Exo-spin™


- **Excellent yield** – Even from very small volumes
- **High exosome purity** – Ultra-low protein and rRNA contamination
- **Consistent results** – Fast and easy protocol

Methods


Exo-spin™ kit can be used to reliably isolate exosomes in 3 easy steps, by combining Precipitation and Size Exclusion Chromatography (SEC), the two most effective methods. A simple representation of this method is shown below.



Results



(A) Size profiles of CSF exosomes isolated using different methods. Data determined by NTA. Each curve represents the average of 3 technical replicate measurements for each exosome isolation method. CSF sample triplicate experiment. (PM = Precipitation Method) (adapted from TS Martins *et al.*, PLoS One 13(6): e0198820 (2018))



(B) Morphology of exosomes pooled by TEM negative staining. Exosomes isolated using Exo-spin™ (adapted from TS Martins *et al.*, PLoS One 13(6): e0198820 (2018)).

For more information on our Exo-spin™ exosome purification kit, please visit our website www.cellgs.com.



Cell Guidance Systems' reagents and services enable control, manipulation and monitoring of the cell, both *in vitro* and *in vivo*

Growth Factors

- Recombinant
- PODS® Sustained Release

Exosomes

- Purification
- Detection
- Purified Exosomes
- NTA Service

Cytogenetics

- Karyotype Analysis
- Array Hybridization

Other research products

- Matrix Proteins
- Small Molecules
- Cell Counting Reagent
- Lipid Quantification Assay



General info@cellgs.com
Technical Enquiries tech@cellgs.com
Orders order@cellgs.com

www.cellgs.com

EUROPE
Cell Guidance Systems Ltd
Maia Building
Babraham Bioscience Campus
Cambridge
CB22 3AT
United Kingdom
T +44 (0) 1223 967316
F +44 (0) 1223 750186

USA
Cell Guidance Systems LLC
Helix Center
1100 Corporate Square Drive
St. Louis
MO 63132
USA
T 760 450 4304
F 314 485 5424