

## GFH170 Recombinant Human GRO- $\gamma$ / CXCL3

### Description

Growth regulated protein  $\gamma$  (GRO- $\gamma$ ), also called CXCL3, acts through the chemokine receptor CXCR2 to promote monocyte migration and adhesion. GRO- $\gamma$  also induces the migration of cerebellar granule neuron precursor cells.

<b>Length</b>	73 aa
<b>Molecular Weight</b>	7.9 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P19876
<b>Purity</b>	$\geq$ 95% determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	Growth regulated protein $\beta$ , GRO $\beta$ , GRO $\beta$ 1, CXCL2, MGSA $\beta$ , MGSA $\beta$ , MGSA-b, CXCL2, MIP-2 $\alpha$ , MIP-2a, MIP2- $\alpha$ , MIP2A, MIP22, GRO2, chemokine (C-X-C motif) ligand 2, C-X-C motif chemokine 2, SCYB2, melanoma growth stimulatory activity $\beta$ , CINC-2a, GROB3, GRO2 oncogene, macrophage inflammatory protein 2 $\alpha$
<b>Biological Activity</b>	Human GRO- $\gamma$ is fully biologically active when compared to standard. The activity is determined by the induced chemotaxis of human neutrophils.
<b>Endotoxin Level</b>	$\leq$ 1.00 EU/ $\mu$ g as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
<b>AA Sequence</b>	ASVVTELRCQ CIQTLOGIHL KNIQSVNVRS PGPHCAQTEV IATLKNGKKA CLNPASPMVQ KII EKILNKG STN


### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

### Data



Induced chemotaxis of human neutrophils assay for Human GRO- $\gamma$ .



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1  $\mu$ g of protein was loaded in each lane. Human GRO- $\gamma$  has a predicted Mw of 7.9 kDa.