

GFM60 Recombinant Mouse IL-17E / IL-25

Description

Interleukin-17E (IL-17E), also known as IL-25, is a member of the IL-17 family of cytokines. IL-17E binds to the IL-17BR receptor to stimulate the secretion of the proinflammatory Interleukin-8 (IL-8) chemokine and to induce the activation of nuclear factor kappa-light-chain-enhancer of activated B cells (NF-kappaB).

Length	154 / 308 aa
Molecular Weight	17.6 / 35.2 kDa
Source	E. coli
Accession Number	Q8VHC9
Purity	≥90% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Interleukin-17E, interleukin 17E, IL17E, IL 17E, Interleukin-25, interleukin 25, IL25, IL 25, IL-25
Biological Activity	Mouse IL-17E is fully biologically active when compared to standard. The activity is determined by the ability to produce human IL-8 by human PBMC cells.
Endotoxin Level	≤1.00 EU/μg as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	MVSLRIQEGC SHLPSCCPK EQEPPEEWLK WSSASVSPPE PLSHTHHAES CRASKDGPLN SRAISPWSYE LDRDLNRVPQ DLYHARCLCP HCVSLQTGSH MDPLGNSVPL YHNQTVFYRR PCHGEEGTHR RYCLERRLYR VSLACVCVRP RVMA

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile 10 mM HCl at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Induced IL-8 production by human PBMC cells when cultured with Mouse IL-17E.