

GFM89AF Recombinant Mouse RELM- γ (Animal-Free)

Description

Resistin-like molecule- γ (RELM- γ) is a member of the RELM-family of secreted proteins containing C-terminal cysteines. The RELM-family consists of resistin (FIZZ3), RELM- α (FIZZ1), RELM- β (FIZZ2), and RELM- γ (FIZZ4). RELM- γ is secreted by peripheral blood granulocytes, bone marrow, spleen, intestine, and lung. RELM- γ functions to promote and regulate promyelocytic differentiation, in addition to regulating nutrient-associated insulin sensitivity in the intestinal tract. Rodents secrete all four RELM-family members, whereas resistin and RELM- β are the only RELM-family members found in humans.

This product is produced with no animal derived raw products. All processing and handling employs animal free equipment and animal free protocols.

Length	89 / 178 aa
Molecular Weight	9.4 / 18.9 kDa
Source	E. coli
Accession Number	Q7TM98
Purity	$\geq 95\%$ determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Resistin-like molecule- γ , RELM γ , FIZZ4
Biological Activity	There is no biological assay data available but protein content is validated by western blot.
Endotoxin Level	≤ 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5
AA Sequence	MEGTLESIVE KVKKELLANR DDCPSTVTKT FSCTSITASG RLASCPSGMT VTGCACGYGC GSWDIRDGNT CHCQCSTMDW ATARCCQLA

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μ g of protein was loaded in each lane. Mouse RELM- γ has a predicted Mw of 18.9 kDa (each monomer is 9.4 kDa). The dimer runs higher in the gel.