


Protocol and instructions

Karyotype Service

**Fixed cells for analysis
from suspension cell culture**

Cat K01, and K05



Contents

Preparing cells	3
A. Reagent preparation for suspension cell culture	3
B. Cell requirements for suspension cell culture	3
C. Procedure for suspension cell culture	4
Sample details	5
A. Requisition form	5
Shipping instructions.....	5
A. Shipping hazardous materials.....	5
B. Prepare the documents.....	5
C. Ship the package	6
D. Packaging Marks	7
E. Delivery address	8

A guide to preparing and shipping fixed cells for karyotype analysis

The results we have seen from labs performing this procedure are quite variable. Be careful to perform the steps as described. We provide a full processing service from live cells. Please consider this as an option.

Whilst we will endeavour to characterise 20 cells (for stable, non-cancerous samples) for poor preparations, finding mitotic figures becomes very time consuming and a full charge will apply if any cells can be analyzed.

Please do not send any samples without contacting us and booking a specific slot.

Preparing cells

This protocol is for preparing cell cultures for karyotype analysis.

Multiple samples can be prepared in parallel using this protocol, however, we recommend fixing no more than 3 x T25 flasks at any one time.

A. Reagent preparation for suspension cell culture

- Pre-made 0.075 M KCl solution (e.g. Thermo Fisher Scientific, cat 10575090) is recommended

This solution should be warmed to 37°C prior to use.

- **Freshly prepared** 3:1 Fixative (add 3 volumes of methanol to 1 volume acetic acid). Methanol (e.g. VWR, Cat 20847.307); Acetic Acid (e.g. VWR, cat 20103.330)

Fixative solution should be prepared on the day and chilled in the fridge (4°C) prior to use.

B. Cell requirements for suspension cell culture

- For each sample, one T25 flask (or equivalent) is required.



Culture should be in log phase, undergoing active cell division. Non-log scale cultures will not yield sufficient cells for analysis. Cultures growing as a monolayer should be at least 50% confluent. Media must be changed 16 – 24 hours prior to harvesting to stimulate cell division.

The following procedure is for T25 flasks (25 cm² cell growth surface). Any reagent volumes should be adjusted according to the cell growth surface used.

C. Procedure for suspension cell culture

1. Adjust media volume to 6 ml.
2. Whilst the suspension cells are in culture, take a representative sample of the growth media/cells from the flask, and use this to perform a cell count. If the cell count shows approximately 70 – 90% of the flask's capacity of cells, then assume the culture is in log phase and proceed with the protocol.
3. Add 60 µl colcemid (10 µg/ml; e.g. Thermo Fisher Scientific, cat 15210040). (If needed, a different volume of media may be used. Add 10 µl of colcemid for every ml of media).
4. Incubate at 37°C for 30 minutes (ESC cultures) or 60 minutes (iPSC cultures). (Please contact us for colcemid incubation times for other cell types).
5. Transfer treated, cell-containing media from flask/dish to a 15 ml centrifuge tube containing 500 µl FBS.
6. Centrifuge the tube at 800 – 1000 rpm (approx. 150 x g) for 8 minutes.
7. Discard the supernatant and flick tube 20 times to dislodge the cell pellet.
8. Using a dropper or a 2 ml serological pipette add 2 ml pre-warmed 0.075 M KCl drop-by-drop (1 drop/second), gently mix. Adjust total volume to 4 ml with 0.075 M KCl in the same manner as before and gently mix again.
9. Incubate at 37°C for 25 minutes.

Note: After hypotonic treatment, cells are more fragile. Handle carefully.

10. Using a dropper or a 2 ml serological pipette, **very slowly** (1 drop/second), add 10 drops of cold fixative (3:1 methanol/acetic acid). Tilt the centrifuge tube and make sure the 10 drops of fixative slide down the side of the tube into the mixture. Mix by gently inverting once or twice.
11. Incubate at room temperature for 10 minutes.
12. Centrifuge the tube at 800 – 1000 rpm (approx. 150 x g) for 8 minutes.
13. Discard the supernatant and flick tube 20 times to dislodge the cell pellet.
14. **Very slowly** (1 drop/second) with gentle mixing between each 3 drops, add 2 ml cold fixative, invert to mix. Adjust the volume of fixative drop by drop (1 drop/second) with gentle mixing to 4 ml, invert to mix.
15. Incubate sample at room temperature for 30 minutes.
16. Centrifuge the tube at 800 – 1000 rpm (approx. 150 x g) for 8 minutes.
17. Discard the supernatant and flick tube 20 times to dislodge the cell pellet.
18. Add 1.5 ml fixative. Swirl to mix.
19. Transfer the cell suspension to a 2 ml screw cap microcentrifuge tube and seal with parafilm.

Please do not use Cryovials as these cause the liquid to splash upon opening and can lead to significant sample loss.

Sample details

A. Requisition form

Please complete a requisition form for each batch of samples being sent to Cell Guidance Systems in advance. The requisition form should be filled and submitted [here](#).

Shipping instructions

A. Shipping hazardous materials

The cell fixative used to prepare cells for shipping is a 3:1 ratio of methanol:acetic acid. Both solutions are considered hazardous materials. Cells suspended in this fixative are non-viable/non-infectious and can be shipped by air in limited (excepted) quantities.

Packaging must be in compliance with the harmonized Excepted Quantity provision of the International Transport Association (IATA), International Civil Aviation Organization (ICAO), and the U.S. Department of Transportation (DOT) regulations.

Quantity limits

Cell pellets can be shipped under the Excepted Quantity provision of IATA in no more than 2 ml of fixative per tube (inner packaging unit) and no more than 50 ml of combined fixative volume per box (outer packing unit).

For example, one shipping box can accommodate a maximum of 25 cell pellets (each cell pellet re-suspended in no more than 1.5 – 2.0 ml of fixative).

Dangerous Goods	UN Identifier	Hazard Class	Packaging Group	Excepted amount per inner package	Excepted amount per outer package
Methanol (flammable)	UN1230	3	2	2 ml	50 ml
Glacial acetic acid (corrosive)	UN2789	8	2	2 ml	50 ml

B. Prepare the documents

The instructions provided below are for FedEx, but you can select your preferred courier provider.

1. Complete a FedEx Expanded Service International Air Waybill. Contact your local FedEx office to assist you in completing this form.

Section	Section Heading	Information to Provide
3	Shipment Information	- Non-viable cells in methanol & acetic acid solution - For medical research use only - Dangerous Goods in Excepted Quantity - IATA 2.7
3	Harmonized Tariff Code	Not applicable

4a	Express Packaging Service	Select: FedEx International Priority
6	Special Handling	Select: Yes, Shipper's Declaration not required

2. Prepare a Commercial Invoice and make 4 copies.

<http://images.fedex.com/downloads/shared/shipdocuments/blankforms/commercialinvoice.pdf>

Section Heading	Information to Provide
Full Description of Goods	<ul style="list-style-type: none"> - Non-viable cells in methanol & acetic acid solution, net 2.0 ml - For medical research use only - Non-infectious, non-hazardous, non-toxic - Dangerous goods in Excepted Quantity - IATA 2.7
Harmonized Code	Not Required
Declaration Code	Not Required

3. E-mail the tracking number and shipment delivery date to Cell Guidance Systems at info@cellgs.com.

C. Ship the package

Packaging for cell pellets fixed in methanol/acetic acid must be constructed with 3 layers:


1. **Inner (primary) packaging.** Re-suspend the cell pellet in 1.5 ml of fixative and place it in a labeled 2 ml microcentrifuge tube. Tighten the cap and seal with parafilm. Do not fill the tube to the top. Leave room for liquid expansion.
2. **Intermediate (secondary) packaging.** Wrap the microcentrifuge tube in absorbent paper toweling and place the tube in a larger 50 ml centrifuge tube. Tighten the cap and seal with parafilm.
3. **Outer packaging.** Place the 50 ml tube in a sturdy, well-padded, cardboard box. Do not send samples in envelopes or similar unsuitable packaging. The dimension of the outer box must be at least 100 mm on two of the three sides.
4. Place the following documents in the shipping box and seal it with packing tape.
 - 1 copy of the Commercial Invoice (EU customers do not require this)
 - MSDS Sheet for glacial acetic acid
https://uk.vwr.com/assetsvc/asset/en_GB/id/7669226/contents
 - MSDS Sheet for methanol
https://uk.vwr.com/assetsvc/asset/en_GB/id/7668343/contents
5. Place 3 copies of the Commercial Invoice on the outside of the box, in the plastic sleeve, under the FedEx Expanded Service International Air Waybill

6. Contact your courier to pick up the box, or arrange to deliver the package to a FedEx drop off point.
7. Email the tracking number and shipment delivery date to Cell Guidance Systems.

D. Packaging Marks

1. Print an Excepted Quantity Package Mark from the attachment and attach it to the outside of the package.
2. Write the name of the sender in the area designated.

**Excepted Quantity Package Mark
From IATA DG Regulations, 50th Ed., Figure 2.7B**



Minimum dimensions of this label are: 100 x 100 mm.

CUT OUT AND ATTACH TO EXTERIOR OF PARCEL.

Hatched image can be in RED or BLACK

Below the hatched E marking record:

Name and address of shipper

DISCLAIMER: These materials are provided as a courtesy, to be used as guidelines to assist properly trained shippers. Cell Guidance Systems is not responsible for correct shipping.

E. Delivery address

For USA

Attention: Umesh Patel
Cell Guidance Systems LLC
Helix Center
1100 Corporate Square Drive
St. Louis
MO 63132
USA
Tel 760 450 4304

For Europe and Rest of the World

Attention: Michael Jones
Cell Guidance Systems
Maia Building
Babraham Bioscience Campus
Cambridge
CB22 3AT
United Kingdom
Tel +44 (0) 1223 497 115

Cell Guidance Systems' reagents and services enable control, manipulation and monitoring of the cell, both *in vitro* and *in vivo*.

Growth Factors

- Recombinant
- Sustained Release

Exosomes

- Purification
- Detection
- Tracking
- NTA Service

Small Molecules

Cell Counting Reagent

Matrix Proteins

Cell Culture Media

- Pluripotent Stem Cells
- Photostable
- *In Vitro* Blastocyst Culture
- ETS-embryo Culture
- Custom Manufacturing Service

Gene Knock-Up System

Cytogenetics Analysis



General info@cellgs.com
Technical Enquiries tech@cellgs.com
Quotes quotes@cellgs.com
Orders order@cellgs.com

www.cellgs.com

EUROPE
Cell Guidance Systems Ltd
Maia Building
Babraham Bioscience Campus
Cambridge
CB22 3AT
United Kingdom
T +44 (0) 1223 967316
F +44 (0) 1223 750186

USA
Cell Guidance Systems LLC
Helix Center
1100 Corporate Square Drive
St. Louis
MO 63132
USA
T 760 450 4304
F 314 485 5424